

University of Arkansas, Fayetteville
ScholarWorks@UARK

Agricultural Economics and Agribusiness
Undergraduate Honors Theses

Agricultural Economics and Agribusiness

5-2017

Cost Benefit Analysis of Genetic Markers in Cattle

Josh Crystal

Follow this and additional works at: <http://scholarworks.uark.edu/aeabuht>



Part of the [Agribusiness Commons](#), [Agricultural Economics Commons](#), and the [Genetics Commons](#)

Recommended Citation

Crystal, Josh, "Cost Benefit Analysis of Genetic Markers in Cattle" (2017). *Agricultural Economics and Agribusiness Undergraduate Honors Theses*. 5.
<http://scholarworks.uark.edu/aeabuht/5>

This Thesis is brought to you for free and open access by the Agricultural Economics and Agribusiness at ScholarWorks@UARK. It has been accepted for inclusion in Agricultural Economics and Agribusiness Undergraduate Honors Theses by an authorized administrator of ScholarWorks@UARK. For more information, please contact ccmiddle@uark.edu, drowens@uark.edu, scholar@uark.edu.

Cost Benefit Analysis on Genetic Markers in Cattle

Josh Crystal

Mentor: Dr. Michael Popp

University of Arkansas

Table of Contents

List of Tables and Figures.....	ii
Abstract.....	iii
Introduction.....	1
Literature Review.....	2
Materials and Methods.....	4
Results.....	10
Discussion.....	11
Conclusion.....	13
References.....	15
Technical Appendix.....	25

List of Tables and Figures

Table 1. Frequencies of Observations by Number of Times Bred for Bermuda and Fescue Pasture Systems, Booneville, AR, 1991-1997.....	19
Table 2. Prices and costs used by FORCAP.....	20
Table 3. Sample summary of cattle and hay management practices on Fescue	21
Table 4. Estimated capital requirements for an operations with 83 cows exposed to a herd sire on 400 acres of continuously grazed pasture with 125 hay acres as modeled in FORCAP	22
Table 5. Multivariate Regression Statistics for Forage Production, Breed, and Marker Effects.....	23
Table 6. Observed and Predicted Profitability in \$/hd by Pasture, Breed, and Marker Effects including seasonality (S+), excluding seasonality (S-), and including ownership charges (S+OC)	24
Appendix Table A1. Analysis of Variance Test for Equality of Means, Observed Means and Estimated Std. Errors for Net Cash Returns (NR) in \$/hd including Seasonality of Sale Prices by Tall Fescue vs. Bermuda Grass Pasture, Cattle Breed (Angus, Brahman and their reciprocal cross) and Genetic Marker, the p450 C994G SNP, Booneville, AR, 1991-97	25
Figure 1. Sample Forage Balance for Fescue (E ⁺) vs. Bermudagrass (BG) Pasture Systems as modeled in FORCAP.....	17
Figure 2. Sample Cattle Input Interface in FORCAP.....	18
Appendix Figure 1. Comparison of Mean Net Returns in \$/hd including Seasonality of Sale Prices by Pasture Based Information (A), Pasture × Breed Based Information (B) and Cross Bred Pasture × Marker Information Based Strategies (C) where Pasture was either Tall Fescue or Bermuda Grass, Cattle Breed was either Angus, Brahman or their reciprocal cross, and the Genetic Marker, the p450 C994G SNP could either have the CC, GC, or GG expression, Booneville, AR, 1991-97. Standard errors are not adjusted for birthweight, breeding failure rate and weaning weight impacts and are different from those estimated using regression analysis.....	26

Abstract

Genetic sequencing in beef cattle (*L. Bos Taurus*) is expected to aid producers select cattle with traits that enhance profitability. Using data from experimental trials conducted with Angus, Brahman, and their reciprocal cross, this project analyzes the profitability of grazing endophyte infected tall Fescue (*L. Festuca arundinacea Schreb.*) vs. Bermuda grass (*L. Cynodon dactylon*) pasture. The study is unique in the sense that actual cow-calf breeding failure rates (open cows were not culled) were tracked from 1991 to 1997 on herds that were bred to calf in spring and either exposed to fungal endophyte (*L. Acremonium coenophialum Morgan-Jones and Gams*) infected tall Fescue grazing and hay or not. Along with data on birth and weaning weights, the number of calves weaned over a cow's useful life were used as major determinants of economic performance using the Forage and Cattle Analysis Planning (FORCAP) decision support software. Using this economic performance measure in conjunction with gene sequencing information allows the selection of beef cattle with profit-maximizing traits (low breeding failure rate) to provide producers with information on which replacement heifers and cows to retain in their breeding stock and herd sires to select. The study examined the single nucleotide polymorphisms (SNPs) p450 C994G, to assess the genetic marker's power to improve the economic performance of the herd. Results suggest that for reciprocal cross herds primarily grazing Bermuda grass pastures the CC expression of the gene is most favorable whereas the GG and GC gene sequences were more profitable with tall Fescue. Adding genetic market information when selecting a production strategy led to approximately \$15/hd in added profitability. At a prorated cost of \$2.40/hd over the life of a dam it is therefore worth pursuing genetic information under the conditions observed in this study.

Introduction

Beef cattle farming has changed drastically over time. It used to be a source of food and power for pulling equipment for a family, but today, more people use cattle farming as a supplemental income source for their family. Larger herd sizes, changing household economics and a better understanding of performance implications of breed selection in light of environmental conditions faced in the field, led to this change. Further, ranchers cross cattle of different breeds to exploit hybrid vigor and improve herd performance by choosing herd sires and replacement heifers with desirable production traits such as low birth weight for calving ease and high weaning weight for greater revenue potential at time of weaning. Genetic selection for lowering breeding failure rate, a stronger indicator of cow herd economics, however, is more difficult and hence scientific advancements capturing an animal's genetic sequence at sufficiently low cost now may allow for decision making using this information.

The objective of this project is to assess whether cow-calf operations would benefit from genetic marker information as they contemplate production practices that would compare the relative profitability of: i) Endophyte infected tall Fescue (E^+) vs. Bermuda grass (BG) pasture management strategies on their operations to assess the impact of presence or absence of fescue toxicosis; ii) the interaction of pasture management with breed selection of purebred Angus, purebred Brahman or their reciprocal cross to demonstrate superior breed selection given available pasture resources; and iii) the interaction of pasture management \times breed \times genetic marker information to determine whether gene sequencing information can help enhance profitability of a cow herd over and above a strategy optimized based on only pasture management and breed information.

Literature Review

Dr. Rosenkrans and other animal scientists have broken down genomes of the bovine by documenting various genetic sequences or genetic markers that make up different phenotypes of cattle often summarized by their expected progeny difference (EPD) that distinguish cattle of a certain breed to a relative moving annual baseline standard either within or across breeds in terms of birth weight, weaning weight, yearling weight, milk, marbling, ribeye area, fat thickness and carcass weight (Kuehn and Thallman, 2016a,b). Hence, farmers examining within- and across-breed based expected progeny differences (EPDs) for making choices involving the genetic makeup of their herd can choose genetics to improve herd performance (Keeton et al., 2014). However, genetic marker information provides a more precise method of developing consistent herd and feedlot performance (Brown et al., 2010; Looper et al., 2010; Rosenkrans et al., 2010; Sales et al., 2011a,b; Thompson et al., 2014).

Looper et al. (2010), conducted a study over the Prolactin (PLR) gene region. This study examined Angus, Brahman, and their reciprocal cross on tall Fescue and Bermuda grasses. They observed the PRL enhancer gene region of the cow to determine if a genetic marker had a noticeable effect on calving rate, birth weight and hip height. The results of the blood test revealed that at two different SNP sites calving rate did not differ (Looper, et al., 2010). This suggests that PLR does aid in mammal reproduction just not as a potential genetic marker. Although, calving ease and birth weights are important for profitability, calving ease and birth weights are not the only traits that can determine profitability in cattle. The study concluded that more research on a larger herd is necessary to determine profitability.

A study conducted by Thompson et al. (2014) observed the value of genetic information for the management and selection of feedlot cattle. The research goal was to determine if cattle possessing differences in their genomes would mature in different ways through the feeding

process. This study looked at 7 traits of feedlot cattle, to determine the expected genetic value of livestock during the feedlot stage (Thompson, et al., 2014). The data obtained was then utilized to estimate differences in average daily gain. Results showed that the benefit cost ratio for selection using certain traits was very low. However, the study did reveal that genetic potential revealed in genetic traits could lead to higher profitability. They found positive changes in profitability of \$22 per head for a single trait and as much as \$38 per head for multiple-traits (Thompson, et al., 2014). This suggests that not only birth weight (BW) but also 205 d weaning weight (WW205) and feedlot performance are important for genetic selection and attendant economic performance.

Sales et al. (2011b) focused on the genetic sequence labeled as p450 C994G among others. This marker is expected to be helpful for determining resistance to toxic fescue. Pastures of E^+ cause a host of detrimental bovine responses. Among them are breeding failures and relatively poor weight gain in offspring as recently analyzed by Caldwell et al. (2013) and Smith et al. (2012). While animal performance is negatively affected by E^+ , pastures of this forage are also more drought tolerant than other cool season forages and many pastures in the mid-Southern U.S. have a significant portion of grazable forage made up of E^+ that is able to persist in pastures despite significant heat stress during the summer months. To combat fescue toxicosis, producers can seed their pastures to BG , free of toxin and heat tolerant, at the cost of added hay feeding during the fall, winter and spring months when cool season fescue would normally offer grazing opportunities for pasture-fed beef cattle (Figure 1). As shown in Figure 1, fescue offers grazing potential primarily in the spring and fall but also part of the summer such that cattle are estimated to graze pasture for 269 d with 96 d on hay using a cow-calf simulation model (Forage and Cattle Analysis and Planning – FORCAP) under conditions that mirrored the E^+ treatment in

Brown et al.'s (1997) study of E^+ and BG pastures. At the same time, grazing of hay land during October and November still allowed 98 bales of excess hay sales given a description of farm conditions detailed below. Cows solely on BG pasture and hay, however, face a different growing season concentrated to summer months and as such graze only 178 d on pasture while being supplemented with hay for 187 d as shown in Figure 1 under the BG treatment conditions using FORCAP where now only 35 excess hay bales could be sold and hayland grazing was not an option as no forage growth occurred during October and November. Hence, using BG leads to more hay feeding but does not expose cattle to deleterious effects of E^+ . Another option for managing E^+ , however, is to modify the genetic makeup of the herd to be more tolerant of these toxins.

Materials and Methods

As described in Brown et al. (1997), purebred Angus, purebred Brahman and their reciprocal cross dams were bred to Hereford sires with calves born in spring of 1991 to 1997 at the Booneville, AR USDA ARS Dale Bumpers Small Farms Research Center (N 35° 5' 42", W 93° 57' 50"). Animals were placed on either E^+ pastures or BG pastures and fed hay of similar type by treatment to ensure a fixed treatment effect of fescue toxicosis that could be observed by breed of dam. To eliminate sire effects, herd sires were rotated across pastures in 13 d intervals throughout the 75 d breeding period. Since some cows had to be replaced in cases where caesarian-sections or uterine prolapse could affect reproductive performance given human interaction or because cows died over the experimental period, not all cows were exposed the same number of times to calculate observed breeding failure rates over the 7 yr study period (Table 1). In addition to observed breeding failure rates defined as:

(1) $BFR = 1 - \# \text{ of calves born} / \# \text{ of times the cow was bred}$

birth weight, sex of calf and 205 d weaning weight data, and genetic marker information on the dam were available along with calving date to capture information needed to perform economic analysis. Cows were fed supplement to ensure adequate body condition score in a similar manner across all treatments.

Using *BFR* along with birth weight, weaning weight, and calving month, the decision support tool for cow-calf operations, FORCAP, was used to capture treatment effects on model farms that primarily differed by pasture system employed as already discussed above (Popp et al., 2013). FORCAP was developed to aid farmers and researchers with analyzing the impacts of:

- input cost and output prices for cattle, fertilizer, feed, marketing charges, fuel, winter annuals, fencing, property tax, insurance, ownership charges of equipment, building and breeding herd investment, operating interest, and major veterinary charges;
- breakdown of land resources for pasture and hay along with amount, type and frequency of fertilizer applications and forage species selection;
- pasture management strategies involving fencing and watering cost, expected grazing efficiency, type and amount of winter annuals, stockpiling and strip grazing preferences;
- cattle production parameters including annual breeding failures, expected cow and calf losses, specification of typical cow replacement age, average mature and young cow weights, weaning age and weights, birth weights and age of first breeding as well as calving season, stocking rate and expected hay waste;
- feeding of extra cattle either retained from own production or purchased using a host supplement feed options;
- managing excess forage using haying equipment, if feasible, in rotationally grazed pastures or fall grazing of hay land when pasture resources are limited.
- hauling and transport decisions
- vaccination program and expected frequency of veterinary services; and
- capital requirements associated with breeding stock, equipment, and buildings along with repair and maintenance, property tax and insurance cost estimates.

In FORCAP, under the ‘Cattle’ tab, a particular cow’s average birth weight for calves as well as their average steer and heifer 205 d weaning weights, *BFR* and average calving month were entered as shown in Figure 2 with the remainder of cow-calf operating parameters held constant except for forage species (Figure 1) utilized. These operating parameters are summarized in Tables 2 to 4 and mirror experimental conditions employed in the original study except that poultry litter was used instead of commercial fertilizer and supplement was not fed as, according to FORCAP, minimal additional nutrition aside from hay was needed to maintain adequate body condition score. As such, net cash return estimates per cow exposed to a herd sire are the revenue from the sale of cattle and excess hay less cash expenses for feed, fertilizer, veterinary and medicine, fuel, twine, repair and maintenance, and operating interest. These net cash returns form the FORCAP estimate of the profitability of a particular animal. Further, it is assumed that the performance of a cow could be replicated or would be similar for a cow with the same genetic marker, breed and pasture management and as such, the profitability estimate of a cow could be extrapolated to herd performance (83 cows in this case to capture a continuously grazed cattle operation for this production region deemed of adequate size to consider obtaining genetic marker information). A ten-year average was used for prices of cattle and fertilizer to remove potential distortion of profitability due to cyclically high or low prices. Seasonality in prices was captured by modifying the calving month and using weaning weight dependent sales prices for the attendant sale month (AMS, 2017). Ten-year average monthly prices for cattle of different weight (Table 2) were also deflated using U.S. All Beef Cattle prices for animals marketed at weights greater than 500 lbs to convert beef prices to constant 2016 dollars (NASS, 2017a). For fertilizer, a fertilizer price index was used to deflate to 2016 dollars (NASS, 2017b).

Finally cost of production estimates for fuel, twine and other inputs were obtained from local sources and reflect cost conditions faced by beef producers in 2016.

Estimates of cow profitability were then regressed against explanatory factors involving genetic marker information, breed, forage species selection, *BFR*, birth and weaning weight variables and select interactions to assess their relative economic impact as follows:

$$(2) \quad NR = a_0 + a_1 \cdot E^+ + a_2 \cdot ANGUS + a_3 \cdot BRAHMAN + a_4 \cdot BFR + a_5 \cdot BW + a_6 \cdot WW205 \\ + a_7 \cdot GC + a_8 \cdot GG + a_9 \cdot E^+ \times ANGUS + a_{10} \cdot BFR \times E^+ + a_{11} \cdot BFR \times ANGUS \\ + a_{12} \cdot BFR \times BRAHMAN + a_{13} \cdot BFR \times GC + a_{14} \cdot BFR \times GG$$

where E^+ is a binary 0/1 variable indicating presence (1) or absence (0) of fescue toxicosis as implied by either using E^+ as the primary pasture and hay forage species or alternatively using *BG* for pasture and hay signaling the absence of fescue toxicosis, *ANGUS* or *BRAHMAN* are binary variables indicating breed, *GC* and *GG* are binary variables indicating presence or absence of p450 G994C (GC) or p450G994G (GG) marker expressions, *BW* is the average birthweight of steers and heifers born to a particular cow and *WW205* is the 205 d adjusted weaning weight equally weighted between steers and heifers as observed in the experiment. The base line represents a reciprocal cross of Angus \times Brahman with a CC marker expression on *BG* pasture and hay devoid of fescue toxicosis pressure as those observations were most frequent in the data set. *BW* and *WW205* are included as explanatory variables as bull EPD statistics typically include values for these parameters. The error term for Eq. 2 is purposely omitted since estimates of *NR* were obtained deterministically in FORCAP. Finally, alternative specifications of the Eq. 2 were estimated with the final specification chosen to i) capture mainly *BFR* interactions with pasture, breed and marker, but ii) also on the basis of goodness of fit using adjusted R^2 and by evaluating the addition or removal of explanatory interactions of variables,

that warranted inclusion given expert opinion, on the basis of the absolute value of their t-statistic ($|t\text{-stat}| > 1.0$). The latter approach limits effects of multicollinearity on coefficient estimates for explanatory variables (Gujarati, 2007).

To answer the questions raised in the objectives above, *NR* could be calculated and compared across pasture system, breed, and genetic marker. As an example, for the *BG* pasture system with reciprocal cross cattle and the CC marker, or the baseline, using the estimated coefficients in Eq. 2 would lead to a profitability estimate for those type of cattle on *BG* pasture as follows:

$$(3) \quad NR_{BG, Cross, CC} = a_0 + a_4 \cdot \overline{BFR}_{BG, Cross, CC} + a_5 \cdot \overline{BW}_{BG, Cross, CC} + a_6 \cdot \overline{WW205}_{BG, Cross, CC}$$

where the a 's are coefficient estimates from Eq. 2 and \overline{BFR} , \overline{BW} , and $\overline{WW205}$ are sub sample averages from observations pertaining to those observations collected on *BG* pastures for reciprocal cross cattle with the CC marker. Changing to E^+ pastures for cattle of the same breed and marker, the applicable additional coefficients, a_1 and a_{10} as shown in Eq. 2, were used with subsample averages for *BFR*, *BW* and *WW205* for cattle on E^+ in a similar fashion as shown in Eq. 3. This process was repeated for the 2 pastures \times 3 breeds \times 3 marker expressions for estimates of 18 *NR* for each treatment combination. To allow comparisons of *NR* across pasture and pasture \times breed, the technical appendix highlights analysis of variance (ANOVA) results undertaken for those comparisons. Further, As the ANOVA did not use explanatory variables of *BFR*, *BW*, and *WW205*, the technical appendix provides the pasture \times breed \times marker comparisons for illustration purposes only.

Ultimately, the target was to analyze if the *NR* of the best marker based selection for a particular forage and breed exceeded the pasture based and/or the pasture \times breed based selection by more than the cost of getting the marker information to then determine if the cow-calf

operator would benefit from getting herd sires and his/her existing breeding stock sampled for the p450 C994G marker.

In order to have a cow tested for genetic markers you must either collect hair or draw blood from the animal. The producer would incur negligible cost to collect hair whereas a blood sample is estimated to cost \$3/hd. Once blood or hair is collected there is an additional cost of \$8/hd to have the blood or hair tested. Adding administrative overhead of \$1/hd, a \$12/hd cost would be a conservative estimate for obtaining genetic marker information. Having your herd tested thus represents more than a trivial cost. Note, however, that this test only needs to be performed once over the life of the cow. For the observed study conditions in Brown et al. (1997), the average number of breeding exposures per cow was five. Hence, an annual cost of testing per head would be \$12/5 or \$2.40/hd. Profitability gains with breeding stock selection based on breed \times pasture \times genetic markers compared to breed and breed \times pasture selection thus needs to exceed \$2.40/hd for cow-calf operations analyzed in this study.

Finally, to test for the impact of seasonality and the potential impact of modified levels in breeding stock investment, the net return estimates in Eq. 2 were also calculated ignoring seasonality by setting the calving month to 100% March in Figure 2 for all observations and by including fixed costs for a cow-calf operation by including taxes, insurance and capital recovery charges for the buildings and equipment as listed in Table 4 along with an opportunity cost of capital invested in breeding stock. Regression analyses on *NR* thus were conducted using Net Cash Returns including seasonality as determined by calving month (*S+*), net cash returns without seasonality (*S-*), and, finally, net returns above cash expenses and ownership charges (*S+OC*). Should analyses of *NR* specified in these three different ways lead to the same conclusions, the results are deemed more robust.

Results

This study examined 86 cows. There were a total of 24 Angus cows, 28 Brahman cows, and 34 reciprocal crosses. Of all cows, 37 cows were on E^+ pasture and hay whereas 49 cows grazed on BG pasture and were fed BG hay. The 86 observations were then broken down further into their genetic marker determined from the p450 C499G SNP. There are 3 different genetic marker expressions, CC, GC, GG. Each marker was represented on E^+ (17 CC, 15 GC, 5 GG) and BG (25 CC, 18 GC, 6 GG) with difference across breed and pasture system detailed below.

Regression results breaking down the effects of breeding failure rate, birth weight, weaning weight, breed, marker, and pasture system (Eq. 2) are shown in Table 5. Coefficients were of the expected sign and adj. R^2 suggested that misspecification was not an issue. Further, coefficient estimates were statistically significant and justified estimation of profitability by pasture \times breed \times marker combination. Results are only shown for the regression statistics on cash net returns using seasonally adjusted prices. Including seasonality and ownership charges in profitability estimates had minimal impact on statistical significance and sign of coefficient estimates and were therefore not shown but are available from the authors upon request.

Table 6 demonstrates actual profitability differences by pasture and pasture \times breed, as well as estimated profitability differences by pasture \times breed \times marker that vary whether seasonally adjusted sale prices were used (S+), sales were all based on October sales given a single March calving month (S-) or whether capital costs associated with retaining additional replacement heifers to account for higher culling rates with greater BFR were included along with seasonal sale price effects (S+OC). Note that the estimates of profitability were calculated using the appropriate coefficient estimates from Table 5 for Eq. 2 with appropriate subsample averages depending on production strategy employed as shown in the left-most column of Table

6. Using BG pasture with reciprocal cross cattle and CC marker information, for example, net cash returns per head were estimated as:

$$NR_{S^+,BG,Cross,CC} = 120.48 - 810.87 \cdot 0.04 + 1.08 \cdot 79.1 - 0.06 \cdot 562.5 = 139.72 \text{ or}$$

$$a_0 + a_4 \cdot \overline{BFR}_{BG,Cross,CC} + a_5 \cdot \overline{BW}_{BG,Cross,CC} + a_6 \cdot \overline{WW205}_{BG,Cross,CC}$$

and were reported at \$141.91 in Table 6 using higher precision using subsample averages of explanatory variables. Note that the other coefficients are not used as the E^+ , breed, and genetic marker variables were set to zero for this subsample and hence their coefficient estimates did not play a role.

Discussion

As shown in Table 6, when comparing E^+ to BG forage systems with the average weights and average BFR , E^+ forage systems outperform the BG system (also shown in Appendix Figure 1A). Given the presence of fescue toxicosis this is puzzling. However, when examining Figure 1 this is understandable. The advantage of E^+ forage systems is the ample opportunity to graze throughout the year and therefore feeding less hay. If a producer were thus interested in managing fescue toxicosis using the BG system and paid no attention to breed or genetic markers, his or her choice would be to pursue an E^+ system as the cost of feeding hay for nearly half of the year is simply too high to offset the gain in weaning weight observed in this study even though the ANOVA equality of means test showed no statistically significant differences.

If the producer now adds breed selection to his repertoire of decision making, then the optimal solution is to have an E^+ forage system and the reciprocal cross of Angus \times Brahman cattle (Appendix Figure 1B). A logical explanation for this is hybrid vigor. Angus tend to show better reproductive performance with fewer breeding failures while Brahman deliver heavier weaning weights with the reciprocal crosses excelling on both fronts regardless of pasture system

employed (Table 6). Note that on *BG* systems weaning weights are higher with the elimination of fescue toxicosis. However, heavier calves exacerbate the problem of hay feeding as more forage consumption leads to even greater hay needs and higher weight calves also lead to lower price per cwt. A *BG* \times *BRAHMAN* strategy in particular, showed negative cash returns not only because of hay feeding but also poor reproductive performance with high *BFR*. Adding breed information compared to only using pasture system information led to higher returns with reciprocal cross cattle on *E+* pastures having the highest observed *S+* returns among choices based on breed and pasture systems at \$169.64/hd.

Finally, considering the addition of p450 C994G SNP on *E+* forage systems, the optimal solution is to have the GC marker in reciprocal crossed cattle resulting in an estimated \$184.99/hd cash returns given the conditions enumerated above (Table 6). Negligible *BFR* in conjunction with highest weaning weight when compared to a similar system with the GC marker that had similar low *BFR* showed that lighter weaning weights lead to lower cattle revenue. Not having the most frequently observed CC marker enhanced reproductive performance leading to greater estimated returns. Similar to results shown above for the pasture \times breed-based profit-maximizing choice, the *BG* system was inferior to the *E+* system as crossbred cattle with either the GG or GC marker had larger *BFR*. Higher weaning weights across all markers were not sufficient to offset costs associated with elevated *BFR* with *BG* compared to *E+*. Noteworthy, and not taken into consideration here, however, is the future fate of calves in feedlots starting at lower weaning weight due to their exposure to *E+* pastures. Nonetheless, adding marker information would allow the producer to gain approximately \$15 per head per year (\$184.99/hd with *E+*, Cross, GC vs. \$169.64/hd on *E+*, Cross) which is

approximately six times the cost of getting the added information conservatively estimated at \$2.40/hd above.

The CC marker performed best on *BG* pastures without fescue toxicosis stress. This suggested the p450 C994G marker indeed measures cattle's ability to deal with toxicosis stress and more so on the reproductive performance end (*BFR*) than in terms of weaning weight.

Removing seasonality effects led to the same conclusions as just discussed above. Profitability estimates were lower on average as seasonal sale prices in October are typically the lowest. Adding ownership charges of cattle to reflect added cost of carrying replacement heifers with higher *BFR* also led to similar conclusions. This suggests that while a sensitivity analysis on hay prices, that may also play a role, was not undertaken, that the results above are robust to the changes analyzed.

Conclusion

The research conducted set out to prove whether it would be profitable for a producer to have their cattle genetically tested to gain information about the p450 C994G marker sequence for tolerance to fescue toxicosis. For cow-calf operations using breeds of Angus and Brahman cattle and grazing on tall Fescue or Bermuda grass pastures, as is common for large regions in the U.S., the results suggested that the genetic marker analyzed indeed would allow producers to enhance their operation's profitability in comparison to a strategy selection purely based on pasture management and breed alone. The results are therefore similar to Thompson et al. (2014) findings and add to information already reported by Looper et al. (2010) and Sales et al. (2011 a,b).

A limitation of the study is that a mixed pasture system consisting of both *BG* and *E+* pastures was not among treatments. Further, had genetic marker information been collected on

the calves, weaning weight differences could have been analyzed for genetic marker information. Finally, had calves been tracked through the feedlot stage, an overall economic performance for not only cow-calf operations would have been possible. It is possible that a *BG* pasture system or even a mixed *BG* and *E+* pasture system could lead to better industry performance as both would potentially mitigate fescue toxicosis effects not only at the cow-calf but also through the feedlot stage of the industry.

References

- Agricultural Marketing Service. 2017. Arkansas Weighted Average Prices for Feeder Cattle, Cull Cow and Bull prices. USDA-AMS. Washington DC Little Rock, AR office.
- Brown, M.A., A.H. Brown Jr., W.G. Jackson, and J.R. Miesner. 1997. "Genotype \times Environment Interactions in Angus, Brahman, and Reciprocal Cross Cows and Their Calves Grazing Common Bermudagrass and Endophyte-Infected Tall Fescue Pastures". *Journal of Animal Science* 75(4):920-925. Retrieved from <http://0-search.proquest.com.library.uark.edu/docview/218121899?accountid=8361>
- Brown Jr., A., S. Reiter, M. Brown, Z. Johnson, I. Nabhan, M. Lamb, A. Starnes and C. Rosenkrans Jr. 2010. "Effects of Heat Shock Protein-70 Gene and Forage System on Milk Yield and Composition of Beef Cattle". *The Professional Animal Scientist* 26(4):398-403. [http://dx.doi.org/10.15232/S1080-7446\(15\)30621-5](http://dx.doi.org/10.15232/S1080-7446(15)30621-5)
- Gujarati, D. 2007. Basic Econometrics. New York, NY: McGraw-Hill/Irwin.
- Keeton, D., M. Popp and A. Smith. 2014. "Economic and Environmental Repercussions of Changing Bull Genetics". *Journal of Farm Managers and Rural Appraisers*, 77: 124-144.
- Kuehn, L.A., and R.M. Thallman. 2016a. "Across-Breed EPD Tables for the Year 2016 Adjusted to Breed Differences for Birth Year of 2014." As presented at the 2014 Beef Improvement Federation Conference. Available at <http://www.bifconference.com/bif2016/documents/2016-ABEPD-press-release.pdf>
- Kuehn, L.A., and R.M. Thallman. 2016b. "Mean EPDs reported by different breeds." In: *Proceedings of the Beef Improvement Federation 48th Annual Research Symposium & Meeting*. p. 122-126.
- Caldwell, J., K. Coffey, J. Jennings, D. Philipp, A. Young, J. Tucker, D. Hubbell, T. Hess, M. Looper, C. West, M. Savin, M. Popp, D. Kreider, D. Hallford, and C. Rosenkrans Jr. 2013. "Performance by Spring and Fall-Calving Cows Grazing with Full, Limited, or No Access to Toxic *Neotyphodium Coenophialum*-Infected Tall Fescue. *Journal of Animal Science* 91(1):465-476. doi:10.2527/jas.2011-4603
- Looper, M., S. Black, S. Reiter, R. Okimoto, Z. Johnson, M. Brown, and C. Rosenkrans Jr. 2010. Identification of polymorphisms in the enhancer region of the bovine prolactin gene and association with profitability traits of beef Cattle1. *Professional Animal Scientist* 26(1): 103-108. <http://0-search.proquest.com.library.uark.edu/docview/230411596?accountid=8361>

- National Agricultural Statistics Service. 2017a. U.S. Beef Cattle Prices [Database.]. USDA-NASS, Washington, DC.
https://www.nass.usda.gov/Charts_and_Maps/graphics/data/priceca.txt (accessed 27 Jan, 2017).
- National Agricultural Statistics Service. 2017b. Quick Stats: Fertilizer Totals—Index for price paid, 2011. [Database.]. USDA-NASS, Washington, DC.
<https://quickstats.nass.usda.gov/results/12922365-06BD-36BD-AE5E-AAB63A064677> (accessed 27 Jan, 2017).
- Popp, M., Smith, A., & Keeton, D. (2013). Forage and Cattle Planner (FORCAP). University of Arkansas, Department of Agricultural Economics and Agribusiness.
<http://agribusiness.uark.edu/decision-support-software.php#forcap>
- Rosenkrans Jr., C., A. Banks, S. Reiter, M. Looper. 2010. “Calving traits of crossbred Brahman cows are associated with Heat Shock Protein 70 genetic polymorphisms”. *Animal Reproduction Science* 119(3-4):178-182. doi:10.1016/j.anireprosci.2010.02.005
- Sales, M. K. Murphy, S. Reiter, A. Brown Jr., M. Brown, M. Looper and C. Rosenkrans Jr. 2011a. “Effects of forage type, body condition and single-nucleotide polymorphisms in the bovine cytochrome P450 regulatory region on cow productivity”. *Journal of Animal Physiology and Animal Nutrition* 97(1):91-96. doi: 10.1111/j.1439-0396.2011.01246.x
- Sales, M., M. Larson, S. Reiter, A. Brown Jr., M. Brown, M. Looper, K. Coffey, and C. Rosenkrans Jr. 2011b. “Effects of bovine cytochrome P450 single-nucleotide polymorphism, forage type and body condition on production traits in cattle.” *Journal of Animal Physiology and Animal Nutrition* 96(4):545-553. doi: 10.1111/j.1439-0396.2011.01176.x
- Smith, S.A., J.D. Caldwell, M. Popp, K.P. Coffey, J.A. Jennings, M.C. Savin, and C.F. Rosenkrans, Jr. 2012. “Tall Fescue Toxicosis Mitigation Strategies: Comparisons of Cow-Calf Returns in Spring- and Fall-Calving Herds.” *Journal of Agricultural and Applied Economics* 44(4):577-592. Retrieved from
<http://ageconsearch.umn.edu/bitstream/137129/2/jaae542.pdf>
- Thompson, N.M., E.A. De Vuyst, B.W. Brorsen, and J.L. Lusk. 2014. Value of genetic information for management and selection of feedlot cattle. *Journal of Agricultural and Resource Economics* 39(1):139-155. Retrieved from <http://0-search.proquest.com.library.uark.edu/docview/1526511508?accountid=836>

Figure 1. Sample Forage Balance for Fescue (E⁺) vs. Bermudagrass (BG) Pasture Systems as modeled in FORCAP.






Figure 2. Sample Cattle Input Interface in FORCAP.



U of A
DIVISION OF AGRICULTURE
RESEARCH & EXTENSION
University of Arkansas System

Cattle Herd Make Up and Performance Statistics



FORCAP
FORAGE & CATTLE PLANNER

Press Reset for default values below		Reset	Press OK below for defaults based on 'Your Farm' options or enter your own below and to the right		Your Farm (includes impact of extra cattle)																		
Description	Bench Mark																						
Days on Hay & Supplements	169				96																		
Days on Pasture	196				269																		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>1. Breeding failures 14% <input type="text" value="OK"/> 0%</p> <p>Cow death losses 1.0% <input type="text" value="OK"/> 1.0%</p> <p>Calf death losses 3.0% <input type="text" value="OK"/> 3.0%</p> <p>Avg. culling age of cows 7.83 <input type="text" value="OK"/> 6.83</p> <p>Avg. number of calves over life of cow 6 <input type="text" value="OK"/> 5</p> <p>Weight of mature cow in lbs 1,250 <input type="text" value="OK"/> 1,200</p> <p>Weight of young cow (at first calf) in lbs 1,000 <input type="text" value="OK"/> 900</p> <p>Weaning age in months 7 <input type="text" value="OK"/> 7</p> <p>Avg. age of replacements at first breeding 15 <input type="text" value="OK"/> 15</p> <p>Avg. birth weight in lbs 90 <input type="text" value="OK"/> 90</p> <p>Avg. steer weaning weight in lbs 555 <input type="text" value="OK"/> 555</p> <p>Avg. heifer weaning weight in lbs 520 <input type="text" value="OK"/> 520</p> <p>Avg. herd sire weight in lbs 2,000 <input type="text" value="OK"/> 1,850</p> </div> <div style="width: 45%;"> <p>Replacement 10 ... check breeding failures and cow losses 17</p> <p>Herd Sires 3 ... based on 'Bull Estimator' 4</p> <p>Calves Sold</p> <p>Male 25 ... based on calf losses, 50/50 steer/heifer calf ratio, replacements retained, cow losses and average culling age as listed in the 'Bull Estimator' 40</p> <p>Female (you buy replacements if negative) 15 23</p> <p>Cull cows 9 16</p> <p>No. of years between bull purchases 1.33 1.00</p> <p>Death losses</p> <p>Cows 1 1</p> <p>Calves 2 3</p> <p>Hay Waste with feeding & storage 20% <input type="text" value="OK"/> 20%</p> <p>Hay produced (from hay & pasture acres in bales) 371 355</p> <p>Hay fed (in bales – accounts for waste) 350 257</p> <p>Number of 1200 lb. round bales sold (bought if negative) 21 98</p> <p>Pasture acres per cow 3.0 ... bench mark changes with fertilizer option (Farm), modify on your farm by selecting a different cow herd size 4.8</p> </div> </div>																							
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>Calving Season</p> <p>... choose your calving season Year round Spring</p> <p>OR provide more detail for your farm</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th>Birth Month(s)</th> <th>% of calves born</th> <th>Sale Month(s)</th> </tr> </thead> <tbody> <tr> <td>Jan</td> <td>0%</td> <td>Aug</td> </tr> <tr> <td>Feb</td> <td>25%</td> <td>Sep</td> </tr> <tr> <td>Mar</td> <td>50%</td> <td>Oct</td> </tr> <tr> <td>Apr</td> <td>25%</td> <td>Nov</td> </tr> <tr> <td>Total</td> <td>100%</td> <td></td> </tr> </tbody> </table> </div> <div style="width: 45%;"> <p>4. Net Cash Returns (\$) 15,768 \$189.98</p> <p>5. GHG (lbs. CO₂ / lb sold) 22.59</p> </div> </div>						Birth Month(s)	% of calves born	Sale Month(s)	Jan	0%	Aug	Feb	25%	Sep	Mar	50%	Oct	Apr	25%	Nov	Total	100%	
Birth Month(s)	% of calves born	Sale Month(s)																					
Jan	0%	Aug																					
Feb	25%	Sep																					
Mar	50%	Oct																					
Apr	25%	Nov																					
Total	100%																						

Note: Ignore the bench mark farm information and focus on the Your Farm information that is tailored to a particular cow deemed representative of the herd. Adjusted for each cow were, (1) breeding failure rate to estimate herd performance if all cows had a specific genetic market of the experimental cow, (2) birth and weaning weights observed after 7 months on pasture, (3) modal calving month for the cow to reflect potential for seasonal price fluctuation, (4) estimated farm level net cash returns defined as cattle and excess hay (5) revenue less cash expenses for feed, fertilizer, veterinary and medicine, fuel, twine, repair and maintenance, and operating interest and divided by number of cows (6) exposed to the herd sires.

Table 1. Frequencies of Observations by Number of Times Bred for Bermuda and Fescue Pasture Systems, Booneville, AR, 1991-1997.

Number of times bred¹	Bermuda²	Fescue²
	# of observations	
7	6 ³	6 ³
6	12	6
5	6	7
4	12	4
3	5	6
2	6	6
1	2	2
Total number of cows per pasture system	49	37

Notes:

¹ Number of times a cow was bred over the period 1991-1997.

² Pasture systems on the basis of primary forage of Bermudagrass (BG) or endophyte-infected tall fescue (E⁺).

³ Number of cows per number of times bred. Frequency distributions were not statistically different (p=0.29) across pasture systems.

Table 2. Prices and costs used by FORCAP¹.

Item and Description	Unit	Price	Item and Description	Unit	Price
Livestock			Feed		
4 - 500 lb. steers ²	\$/cwt	170.62	Hay delivered/sold FOB -- 5' x 5' (1,200 lbs)	\$/bale	60.00
5 - 600 lb. steers	\$/cwt	153.48	Salt & minerals (50 lb bag)	\$/bag	20.00
6 - 700 lb. steers	\$/cwt	142.41	Fertilizer		
7 - 800 lb. steers	\$/cwt	136.83	Lime	\$/ton	33.10
3 - 400 lb. heifers	\$/cwt	146.13	Ammonium nitrate (34-0-0)	\$/ton	338.64
4 - 500 lb. heifers	\$/cwt	135.48	Poultry litter (3-2-3)	\$/ton	18.74
5 - 600 lb. heifers	\$/cwt	129.15	Application cost per acre	\$/acre	4.61
6 - 700 lb. heifers	\$/cwt	125.76	Fuel Use & Other Miscellaneous		
Cull cow ³	\$/cwt	64.35	Amortized pasture/hay maintenance & establishment ⁵	\$/acre	14.00
Purchase price of breeding bull	\$/hd	2,000	Fuel use for mowing, raking, and staging	gal/acre	4.50
Cull bull ⁴	\$/cwt	80.77	Fuel use per day for feeding	gal/70 cows/day	1.19
Beef check off, ins. & yardage	\$/hd	1.00	Fuel use per day for checking cattle	gal	1.00
Sales commission (% of sales)	%	3.50	Fuel cost	\$/gal	1.70
Veterinary Services Charges			Twine	\$/bale	1.00
Prolapse	\$/hd	75	Cost for farm vehicle (\$/hd/month)	\$	1.00
Caesarian section	\$/hd	225	Capital recovery rate ⁶	%	5.00
Sick treatment (avg. drug charge)	\$/hd	15	Operating interest ⁷	%	4.75
Bull soundness	\$/hd	30			

Notes:

¹ Forage and Cattle Planner.

² State average, medium and large frame No. 1 prices as reported by the United States Department of Agriculture, Agricultural Market Service. A ten year average was used for sale months that were split across several marketing months with a specific calving distribution and depended on weaning age. Shown are the sale prices for cattle when selecting a spring calving season with 25% of calves born in February and April and 50% born in March. Prices were deflated using average US beef cattle prices. Further, calf prices are linearly interpolated across weight categories to adjust for specific sale weight.

³ 75-80% Lean Breaking Utility.

⁴ Yield Grade 1-2, 1,000 to 2,100 lbs.

⁵ Based on 10-year life of stand and standard seedbed preparation and weed control expenses.

⁶ Capital recovery rate is used for estimating ownership charges on equipment and buildings and is also used for the opportunity cost of investment in breeding stock.

⁷ Charged on half the cash operating expenses incurred per year to reflect likely operating credit line expense.

Table 3. Sample¹ summary of cattle and hay management practices on Fescue.

Description		Herd Size and Description	
Days on hay & supplements ²	96	Cows (avg. age 65 months)	66
Days on pasture ²	269	Young cows (avg. age 30 months)	17
Breeding failures ³	18%	Cow herd size ⁶	83
Annual cow death losses	1%	Replacement heifers	17
Annual calf death losses	3%	Herd sires or bulls needed ⁷	4
Avg. culling age of cows	6.83	Male calves sold	33
Avg. number of calves over life of cow ⁴	5	Female calves sold	16
Weight of mature cow in lbs	1,200	Cull cows	16
Weight of young cow (at first calf) in lbs	900	No. of years between bull purchases	1.00
Weaning age in months ⁵	7	Cow death losses	1
Avg. age at first breeding	15	Calf death losses	2
Avg. birth weight in lbs	80	Hay waste with feeding & storage	20%
Avg. steer weaning weight in lbs	492	Hay (1,200 lb. round bales)⁸	
Avg. heifer weaning weight in lbs	464	Hay produced	355
Avg. bull weight in lbs	1,850	Hay sold (purchased)	98
Calving season	User-selected ⁵	Pasture acres per cow ⁵	4.8

Notes:

- ¹ Results are for a cow-calf operation with 125 acres of hay and 400 acres of pasture planted to endophyte-infected tall fescue (E⁺). Fertilizer is added in the form of poultry litter (3-2-3) on hay and pasture at a rate of 2 tons per acre. Pastures are continuously grazed and cattle are allowed to graze hayland during October and November.
- ² Will vary with calving month, breeding failure rate and calf weaning weight as grazing demand considers animal weight and herd size on a month to month basis.
- ³ Varies by cow and grazing system. Higher breeding failure rates are expected for E⁺ pastures. Actual experimentally observed long term breeding failure rates are used by cow.
- ⁴ Determines the culling age for cows and hence the number of replacement cows needed in conjunction with breeding failures.
- ⁵ Calving season affects sale prices and seasonal nutrient needs of the herd.
- ⁶ Cow herd makeup is described for first-bred, young cows, and older cows in the rows above the cow herd size row. The cow herd size is the number of animals bred per year. Cow herd size also drives the stocking rate shown in pasture acres per cow.
- ⁷ The number of herd sires is determined by the total number of first-bred, young cows, and older cows in the herd. This analysis assumes that one bull can service a maximum of 25 cows. Bulls are assumed to be replaced every four years.
- ⁸ A negative number implies greater hay needs than available from farm production and hence hay purchases.

Table 4. Estimated capital requirements for an operations with 83 cows exposed to a herd sire on 400 acres of continuously grazed pasture with 125 hay acres as modeled in FORCAP¹.

Description	List Price	Years of Useful Life	Salvage Value	Capital Recovery	Repair Factor	Repair and Maintenance	Insurance & Property Taxes
Hay Barn (1,000 sq. ft.)	\$5,000	20	\$800	\$377	0.40	\$100	\$65
Shed (800 sq. ft.)	\$4,000	20	\$750	\$298	0.40	\$80	\$52
50-75 hp Tractor	\$30,000	10	\$10,000	\$3,090	0.25	\$750	\$390
Disk Mower	\$8,000	7	\$4,000	\$891	0.35	\$400	\$104
Hay Baler	\$20,000	10	\$7,500	\$1,994	0.10	\$200	\$260
Hay Rake	\$4,000	10	\$750	\$458	0.20	\$80	\$52
Stock Trailer	\$3,500	10	\$1,500	\$334	0.20	\$70	\$46
Hay Wagon	\$3,000	10	\$500	\$349	0.20	\$60	\$39
Brush Mower	\$8,000	10	\$800	\$972	0.25	\$200	\$104
Corral and Chute	\$3,500	10	\$1,000	\$374	0.15	\$53	\$46
Miscellaneous Items	\$2,000	10	\$0	\$259	0.50	\$100	\$26
Fencing & Watering	\$33,306	20	\$0	\$2,673	0.10	\$167	-
Total Equipment & Buildings	\$124,306			\$12,069		\$2,259	\$1,183
Breeding Stock	Qty	Price	Value				
Cows	66	\$850	\$56,100	\$2,805			
Young Cows ²	17	\$1,000	\$17,000	\$850			
Replacement Heifers ²	17	\$900	\$15,300	\$765			
Herd Sires	4	\$2,000	\$8,000	\$400			
Total Breeding Stock	104		\$96,400	\$4,820			

Notes:

¹ Forage and Cattle Planner.

² The need for more or fewer replacement heifers given changes in breeding failure rates modifies breeding stock investment and thereby the opportunity cost of capital employed in the cattle operation. Opportunity cost represents returns foregone if money invested in cattle were invested elsewhere. As such, the annual opportunity cost of cattle is \$96,400 times the 5% capital recovery rate.

Table 5. Multivariate Regression Statistics for Forage Production, Breed, and Marker Effects.

Variable		Coefficient (Std. Error)	T-Statistic
Constant	a_0	119.79 (43.14) ^{***,1}	2.78
$E^{+,2}$	a_1	3.39 (9.61)	0.35
<i>ANGUS</i>	a_2	-57.65 (13.00) ^{***}	-4.44
<i>BRAHMAN</i>	a_3	2.49 (11.37)	0.22
<i>BFR</i>	a_4	-808.88 (44.41) ^{***}	-18.21
<i>BW</i>	a_5	1.11 (0.43) ^{**}	2.58
<i>WW205</i>	a_6	-0.06 (0.08)	-0.80
<i>GC</i>	a_7	4.25 (8.46)	0.50
<i>GG</i>	a_8	-5.53 (12.03)	-0.46
$E^+ \times \textit{ANGUS}$	a_9	50.10 (13.77) ^{***}	-3.64
$\textit{BFR} \times E^+$	a_{10}	-156.72 (29.40) ^{**}	5.33
$\textit{BFR} \times \textit{ANGUS}$	a_{11}	144.77 (50.70) ^{***}	2.86
$\textit{BFR} \times \textit{BRAHMAN}$	a_{12}	-53.04 (47.67) ^{**}	-1.11
$\textit{BFR} \times \textit{GC}$	a_{13}	35.06 (32.05)	1.09
$\textit{BFR} \times \textit{GG}$	a_{14}	105.98 (53.25) [*]	1.99
R^2		97.65%	
Adj. R^2		97.19%	
# of obs.		86	

Notes:

¹ * < 0.1, ** < 0.05, and *** < 0.001 level of significance.² E^+ is a binary (0/1) variable and represents the presence of endophyte infected tall fescue as feed source on pasture and from hay. *ANGUS*, *BRAHMAN*, *GC*, and *GG* are also binary variables indicating presence = 1 or absence = 0 of breed and genetic marker *p450 GC* and *p450 GG*, respectively. *BFR*, *BW*, and *WW205* are cow specific average 1991 – 1997 performance statistics related to breeding failure rate, average birth and weaning weight, respectively. The baseline scenario reflects a Bermudagrass (BG) pasture system devoid of fescue toxicosis using reciprocal cross cattle with the *p450 CC* genetic marker expression.

Table 6. Observed and Predicted Profitability in \$/hd by Pasture, Breed, and Marker Effects including seasonality (S+), excluding seasonality (S-), and including ownership charges (S+OC).

Description	# of obs.	Avg. of Explanatory Variables ¹			Profitability in \$/hd ²		
		BW	BFR	WW205	S+	S-	S+OC
<i>E</i> ⁺ ³	37	79.7	16.5%	477.5	\$54.56	\$50.61	-\$173.47
<i>BG</i>	49	79.8	17.8%	546.4	\$19.54	\$5.06	-\$208.07
<i>E</i> ⁺ × <i>ANGUS</i>	10	79.2	18.9%	386.3	-\$6.71	-\$9.15	-\$235.21
<i>E</i> ⁺ × <i>CROSS</i>	15	81.8	2.2%	522.8	\$169.64	\$167.73	-\$54.39
<i>E</i> ⁺ × <i>BRAHMAN</i>	12	77.5	32.5%	496.9	-\$38.24	-\$45.99	-\$270.88
<i>BG</i> × <i>ANGUS</i>	14	83.2	12.1%	488.4	\$49.83	\$36.23	-\$175.94
<i>BG</i> × <i>CROSS</i>	19	78.1	6.8%	571.6	\$119.57	\$101.16	-\$104.92
<i>BG</i> × <i>BRAHMAN</i>	16	78.9	35.9%	567.3	-\$125.73	-\$136.33	-\$358.67
<i>E</i> ⁺ × <i>ANGUS</i> × <i>CC</i>	3	75.0	4.7%	377.0	\$52.46	\$55.22	-\$171.77
<i>E</i> ⁺ × <i>ANGUS</i> × <i>GC</i>	5	81.8	30.0%	395.0	-\$54.81	-\$62.87	-\$286.90
<i>E</i> ⁺ × <i>ANGUS</i> × <i>GG</i>	2	79.0	12.5%	378.5	\$24.81	\$28.56	-\$201.10
<i>E</i> ⁺ × <i>CROSS</i> × <i>CC</i>	7	83.6	4.7%	529.4	\$153.52	\$149.18	-\$71.23
<i>E</i> ⁺ × <i>CROSS</i> × <i>GC</i>	6	80.3	0.0%	528.5	\$184.99	\$181.20	-\$38.73
<i>E</i> ⁺ × <i>CROSS</i> × <i>GG</i>	2	80.0	0.0%	482.5	\$177.62	\$179.37	-\$46.09
<i>E</i> ⁺ × <i>BRAHMAN</i> × <i>CC</i>	7	76.3	24.1%	499.1	\$10.23	\$8.15	-\$218.60
<i>E</i> ⁺ × <i>BRAHMAN</i> × <i>GC</i>	4	81.5	47.0%	485.5	-\$123.61	-\$137.32	-\$362.79
<i>E</i> ⁺ × <i>BRAHMAN</i> × <i>GG</i>	1	70.0	33.0%	527.0	-\$31.43	-\$33.91	-\$261.92
<i>BG</i> × <i>ANGUS</i> × <i>CC</i>	4	83.0	18.8%	480.0	\$1.06	-\$13.32	-\$226.18
<i>BG</i> × <i>ANGUS</i> × <i>GC</i>	9	82.6	10.4%	488.7	\$63.11	\$49.15	-\$162.38
<i>BG</i> × <i>ANGUS</i> × <i>GG</i>	1	90.0	0.0%	519.0	\$125.48	\$118.11	-\$96.93
<i>BG</i> × <i>CROSS</i> × <i>CC</i>	10	79.1	4.0%	562.5	\$141.58	\$122.61	-\$82.16
<i>BG</i> × <i>CROSS</i> × <i>GC</i>	6	75.7	11.7%	576.8	\$83.24	\$66.39	-\$142.35
<i>BG</i> × <i>CROSS</i> × <i>GG</i>	3	79.7	6.7%	591.7	\$120.44	\$107.79	-\$103.47
<i>BG</i> × <i>BRAHMAN</i> × <i>CC</i>	11	80.3	34.7%	577.1	-\$122.45	-\$134.35	-\$354.97
<i>BG</i> × <i>BRAHMAN</i> × <i>GC</i>	3	75.7	38.0%	560.7	-\$137.24	-\$149.54	-\$372.50
<i>BG</i> × <i>BRAHMAN</i> × <i>GG</i>	2	76.0	39.5%	523.5	-\$128.79	-\$140.27	-\$361.89

Note:

¹ Birth weight (*BW* in lbs/hd), breeding failure rate (*BFR* as defined in Eq. 1), and weaning weight (*WW205* in lbs/hd averaged across male and female calves per cow) are reported for subsamples meeting the pasture system, breed, and genetic marker characteristics shown in the left most column.

² Profitability numbers are observed averages for pasture (*E*⁺, *BG*) and pasture × breed interactions.

³ *E*⁺ and *BG* represent the presence of endophyte infected tall fescue and Bermudagrass, respectively as the sole feed source on pasture and from hay. *ANGUS*, *BRAHMAN*, *CROSS*, *GC*, and *GG* are variables indicating breed, reciprocal cross, and presence of genetic markers *p450CC*, *p450 GC* and *p450 GG*.

Technical Appendix

Somewhat similar to the linear regression results shown in Table 5 an analysis of variance was conducted in EViews v9 (IHS Global Inc., Irvine, CA) as shown in Appendix Table A1 and Appendix Figure 1.

Appendix Table A1. Analysis of Variance Test for Equality of Means, Observed Means and Estimated Std. Errors for Net Cash Returns (*NR*) in \$/hd including Seasonality of Sale Prices by Tall Fescue vs. Bermuda Grass Pasture, Cattle Breed (Angus, Brahman and their reciprocal cross) and Genetic Marker, the p450 C994G SNP, Booneville, AR, 1991-97.

Information Used	Treatment	Mean ¹	Std Error	model df	error df	F-value (p-value)
Pasture	Fescue (E ⁺)	19.54	23.66	1	84	1.0318
	Bermuda Grass (BG)	54.56	24.31			0.3126
Pasture × Breed	E ⁺ × Angus (AA)	-6.71	36.31	5	80	11.8439
	E ⁺ × Cross (AB)	169.64	16.01			<.0001
	E ⁺ × Brahman (BB)	-38.24	45.64			
	BG × AA	49.83	27.83			
	BG × AB	119.57	23.91			
	BG × BB	-125.73	42.81			
Pasture × Breed × Marker	E ⁺ × AA × CC	61.87	35.37	17	68	3.6691
	E ⁺ × AA × GC	-61.04	61.40			0.0001
	E ⁺ × AA × GG	26.27	48.34			
	E ⁺ × AB × CC	157.64	33.75			
	E ⁺ × AB × GC	187.82	9.32			
	E ⁺ × AB × GG	157.05	4.05			
	E ⁺ × BB × CC	11.20	44.45			
	E ⁺ × BB × GC	-126.74	109.36			
	E ⁺ × BB × GG	-30.29	na			
	BG × AA × CC	-1.97	70.72			
	BG × AA × GC	63.44	29.21			
	BG × AA × GG	134.5	na			
	BG × AB × CC	135.96	17.45			
	BG × AB × GC	85.66	68.00			
	BG × AB × GG	132.72	56.81			
	BG × BB × CC	-122.04	58.76			
	BG × BB × GC	-134.26	90.68			
	BG × BB × GG	-133.22	71.63			

Notes:

¹ Arithmetic means are reported for net cash returns defined as cattle and hay sales less cash costs of fuel, fertilizer, twine, medicine and vet services, repair and maintenance, as well as operating interest.

² Estimated standard errors to use for mean comparisons. Two categories had a single observation.

Appendix Figure 1. Comparison of Mean Net Returns in \$/hd including Seasonality of Sale Prices by Pasture Based Information (A), Pasture \times Breed Based Information (B) and Cross Bred Pasture \times Marker Information Based Strategies (C) where Pasture was either Tall Fescue or Bermuda Grass, Cattle Breed was either Angus, Brahman or their reciprocal cross, and the Genetic Marker, the p450 C994G SNP could either have the CC, GC, or GG expression, Booneville, AR, 1991-97. Standard errors are not adjusted for birthweight, breeding failure rate and weaning weight impacts and as such are different from those estimated using regression analysis.

